

Quarkonium dissociation in a thermal medium

Jakub Jankowski and David Blaschke

Institute for Theoretical Physics, University of Wrocław, pl Maxa Born'a 9, 50-204 Wrocław, Poland

DOI: <http://dx.doi.org/10.3204/DESY-PROC-2008-xx/jankowski.jakub>

We investigate the Mott effect for heavy quarkonia due to Debye screening of the heavy quark potential in a plasma of massless quarks and antiquarks. The influence of residual color correlation is investigated by coupling the light quark sector to a temporal gauge field driven by the Polyakov loop potential. This leads to an increase of the Mott dissociation temperatures for quarkonia states which stabilizes in particular the excited states, but has marginal effect on the ground states.

1 Introduction

Since the suggestion of J/ψ suppression as a signal of quark-gluon plasma (QGP) formation by Matsui and Satz [1] in 1986 the problem of quarkonium dissociation in hot and dense strongly interacting matter has played a key role for QGP diagnostics in relativistic heavy-ion collision experiments. The original idea was that in a QGP the string tension of the confining potential vanishes and the residual one-gluon exchange interaction undergoes a Debye screening by the color charges of the plasma. When the temperature dependent Debye radius $r_D(T)$ (the inverse of the Debye mass $m_D(T)$) becomes shorter than the Bohr radius of the charmonium ground state (J/ψ) then the Mott effect [2] (bound state dissociation) occurs and the corresponding temperature is $T_{\text{Mott}}^{J/\psi}$. This simple idea grew up to a multifaceted research direction when not only in the first light ion - nucleus collisions at the CERN NA38 experiment, but also in proton - nucleus collisions at Fermilab J/ψ suppression has been found so that there is not only a QGP but also a cold nuclear matter effect on charmonium production, see [3] for a recent review.

If one wants to explore the question of screening in a plasma more in detail then a variety of approaches is available in the literature, from the original Debye-Hückel approach [4] where one can study any vacuum potential (for example the Cornell potential) and see its medium modification, over the thermodynamic Green functions approach to the ab-initio studies of heavy-quark potentials in lattice QCD. With the obtained medium-dependent potentials one can then study the bound state problem by solving the nonrelativistic Schrödinger equation or, more systematically, the thermodynamic T - matrix for quarkonia [5].

On the other hand one may calculate proper correlators directly from lattice QCD and extract from them spectral functions [6]. There is an intriguing disagreement between the Mott temperatures deduced from these spectral functions and those of the potential models: the latter are much smaller than the former! From the lattice data for quarkonium correlators one has extracted $T_{J/\psi}^{\text{Mott}} \approx 1.9T_c$ while in potential model calculations $T_{J/\psi}^{\text{Mott}} \approx 1.2T_c$. This problem has led to the discussion of the proper thermodynamical function to be used as a potential

in the Schrödinger equation. Should it be the free energy or the internal energy? We will not follow this question in the present work, but refer to [3, 7] and references therein.

Here we examine a simple model of screening as derived from one-loop calculations in thermal quantum field theory, and make a small extension of this result by putting the internal fermion lines in a constant temporal gauge field which mimics confining gluon dynamics (Polyakov-loop potential). In our approach the medium is made of plasma of massless quarks, described by the chirally symmetric phase of the Nambu–Jona-Lasinio (NJL) model of QCD [8]. Confinement is implemented in the most simple way by coupling the system to the Polyakov loop variable - resulting in the so called Polakov loop NJL-model (or PNJL model). Recently it has been successfully used to reproduce lattice data [9] or to describe light meson physics at finite temperatures and densities [10].

2 Debye-screening in a PNJL quark plasma

Given the self energy (polarization function) of a boson that mediates the interaction, the screened potential is given by a resummation of one-particle irreducible diagrams ("bubble" resummation = RPA) [11]

$$V_{\text{sc}}(q) = V(q)/[1 + F(0; \mathbf{q})/q^2] , \quad (1)$$

where we take $V(q) = -\frac{4}{3}g^2/q^2$, $q^2 = |\mathbf{q}|^2$ as the unscreened vacuum potential. The longitudinal gluon polarization function $F(0; \mathbf{q}) = -\Pi_{00}(0; \mathbf{q})$ in the finite T case can be calculated within perturbative thermal field theory where it takes the form

$$\Pi_{00}(i\omega_l; \mathbf{q}) = Tg^2 \sum_{n=-\infty}^{\infty} \int \frac{d^3p}{(2\pi)^3} \text{Tr}[\gamma^0 S_{\Phi}(i\omega_n; \mathbf{p}) \gamma^0 S_{\Phi}(i\omega_n - i\omega_l; \mathbf{p} - \mathbf{q})] , \quad (2)$$

where $\omega_l = 2\pi lT$ are the bosonic and $\omega_n = (2n+1)\pi T$ are the fermionic Matsubara frequencies of the imaginary-time formalism. The symbol Tr stands for traces in color, flavor and Dirac spaces. S_{Φ} is the propagator of a massless fermion coupled to the homogeneous static gluon background field φ_3 . Its inverse is given by [9, 10]

$$S_{\Phi}^{-1}(\mathbf{p}; \omega_n) = \gamma \cdot \mathbf{p} + \gamma_0 i\omega_n - \lambda_3 \varphi_3 , \quad (3)$$

where φ_3 is related to the Polyakov loop variable defined by [9]

$$\Phi(T) = \frac{1}{3} \text{Tr}_c(e^{i\beta\lambda_3\varphi_3}) = \frac{1}{3}(1 + 2\cos(\beta\varphi_3)) .$$

The physics of $\Phi(T)$ is governed by the temperature-dependent Polyakov loop potential $\mathcal{U}(\Phi)$, which is fitted to describe the lattice data for the pressure of the pure glue system [9]. After performing the color-, flavor- and Dirac traces and making the fermionic Matsubara summation, we obtain in the static, long wavelength limit

$$\Pi_{00}(0; \mathbf{q}) = \frac{2N_{\text{dof}}g^2}{\pi^2} \int_0^{\infty} dp p^2 \frac{\partial f_{\Phi}}{\partial p} = -\frac{4N_{\text{dof}}g^2}{\pi^2} \int_0^{\infty} dp p f_{\Phi}(p) = -\frac{N_{\text{dof}}g^2T^2}{3} I(\Phi) = -m_D^2(T) . \quad (4)$$

where $m_D(T)$ is the Debye mass, the number of degrees of freedom is $N_{\text{dof}} = N_c N_f = 6$ and $f_{\Phi}(p)$ is quark distribution function [10]. The screened potential is thus

$$V_{\text{sc}}(q) = -4\pi\alpha/[q^2 + m_D^2(T)] . \quad (5)$$

In comparison to the free fermion case [11, 12] the coupling to the Polyakov loop variable $\Phi(T)$ gives rise to a modification of the Debye mass, given by the integral

$$I(\Phi) = \frac{12}{\pi^2} \int_0^\infty dx x \frac{\Phi(1 + 2e^{-x})e^{-x} + e^{-3x}}{1 + 3\Phi(1 + e^{-x})e^{-x} + e^{-3x}}. \quad (6)$$

In the limit of deconfinement ($\Phi = 1$), the case of a massless quark gas is obtained ($I(1) = 1$) while for confinement ($\Phi = 0$) one finds that $I(0) = 1/9$. For the temperature dependence of $\Phi(T)$ we employ in the following chapter the results of a nonlocal PNJL model [13].

3 Variational ansatz and estimation of Mott temperatures

Here we will use the derived potential in the quantum mechanical way to estimate the dissociation temperature. With the trial wave function for the 1S state

$$\psi_{1S}(r; \gamma) = \sqrt{\frac{\gamma^3}{\pi}} \exp(-\gamma r) \quad (7)$$

and the non-relativistic two-body Hamiltonian

$$H = -\frac{\nabla^2}{m_Q} - \frac{\alpha}{r} e^{-m_D(T)r}, \quad (8)$$

where the potential term is the Fourier transform of screened Coulomb potential (5), we obtain the energy functional for the Ritz variational principle

$$E_{1S}(\gamma, T) = \langle \psi_{1S}(\gamma) | H | \psi_{1S}(\gamma) \rangle = \frac{\gamma^2}{m_Q} - \frac{4\alpha\gamma^3}{(m_D(T) + 2\gamma)^2}. \quad (9)$$

Simultaneously satisfying the conditions for the ground state energy $dE_{1S}(\gamma, T)/d\gamma = 0$ and for a vanishing binding energy (Mott effect), $E_{1S}(\gamma, T^{\text{Mott}}) = 0$, provides us with an analytic expression for the critical Debye mass $m_D(T_{1S}^{\text{Mott}}) = 2\gamma$. Using once again the condition $E_{1S}(\gamma, T^{\text{Mott}}) = 0$, results in the Mott condition for the Debye potential [2]

$$r_D(T_{1S}^{\text{Mott}}) = a_0, \quad (10)$$

where $a_0 = 2/(\alpha m_Q) = 1/\sqrt{\varepsilon_0 m_Q}$ is the Bohr radius and $\varepsilon_0 = \alpha^2 m_Q/4$ the binding energy of ground state in the vacuum ($m_D = 0$). Due to the temperature dependence of the Debye mass, we obtain the Mott dissociation temperature in the massless quark gas (for $\Phi = 1$)

$$T^{\text{Mott}} = \sqrt{3\varepsilon_0 m_Q / N_{\text{dof}} / g} = \sqrt{\varepsilon_0 m_Q^3 / (2\pi N_{\text{dof}})}. \quad (11)$$

In Tab. 1 we give the parameter values according to set (i) of Ref. [14]. The Polyakov loop variable contribution considered in previous section affects the Mott temperature in Eq. (10) and gives the following formula

$$T^{\text{Mott}, \Phi} = T^{\text{Mott}} / \sqrt{I(\Phi)}. \quad (12)$$

Table 1: Mott temperatures T^{Mott} ($T^{\text{Mott},\Phi}$) according to Eq. (11) (Eq. (12)) for a massless ideal quark gas (PNJL model). The critical temperature is $T_c = 202$ MeV [13]. The parameters are fixed to reproduce quarkonium states in vacuum as Coulombic bound states [14]. In the charmonium (bottomonium) system the heavy quark mass m_Q is $m_c = 1.94$ GeV ($m_b = 5.1$ GeV) and the ground state binding energy ε_0 is 0.78 GeV (0.75 GeV).

	T^{Mott}/T_c	$T^{\text{Mott},\Phi}/T_c$
J/ψ	1.25	1.37
χ_c	0.83	1.11
ψ'	0.66	0.99
Υ	2.50	2.50
χ_b	1.72	1.73
Υ'	1.28	1.39




Figure 1: Temperature dependence of the Debye screening radius $r_D(T)$ for charmonium (upper panel) and bottomonium (middle panel) with (solid lines) and without (dashed lines) coupling to the Polyakov loop $\Phi(T)$ shown in the lower panel (from [13]). Sequential dissociation of quarkonia states occurs at the Mott temperatures T^{Mott} for which their Bohr radius equals $r_D(T)$.

This means that in the case of confining color correlations ($0 \leq \Phi < 1$) the Debye screening radius is larger than in a free quark gas at the same temperature, so that bound states get stabilized against thermal dissociation by color screening.

The influence of the Polyakov loop variable on the dissociation temperature is summarized in Figure 1 which shows the temperature dependence of the Debye radius $r_D(T) = 1/m_D(T)$ compared to the Bohr radii of the low-lying states of the charmonium and bottomonium family, respectively. Due to its larger mass and smaller Bohr radius, the Υ dissociates at higher temperatures than the J/ψ , where the free quark gas case is almost reached and the Mott temperatures which almost coincide for the cases with and without coupling to the Polyakov

loop field. For the lighter J/ψ there is a noticeable stabilization due to the coupling to the Polyakov loop potential which results in an increase of the Mott dissociation temperature from $1.25 T_c$ to $1.37 T_c$, still more similar to results of nonrelativistic potential models rather than the still higher dissociation temperatures conjectured from the spectral functions deduced from lattice data for heavy quarkonium correlators by the maximum entropy method. For estimating the Mott temperatures of the excited quarkonia states we have employed the scaling of bound state radii: $r_{\chi_c} = 1.5 r_{J/\psi}$, $r_{\psi'} = 2.0 r_{J/\psi}$ as obtained in the Cornell-type potential model [7].

4 Conclusions

We have applied the methods of thermal field theory to estimate the effects of Debye screening on heavy quarkonia bound state formation. In order to account for residual effects of confining color correlations in the deconfined phase, we have used the PNJL model in the evaluation of the one-loop polarization function. As expected, a stabilization of bound states in the vicinity of the critical temperature for $T > T_c$ is obtained. We applied Ritz' variational principle to derive the Mott criterion for bound states of the statically screened Debye potential and obtained Mott temperatures in good agreement with previous results from nonrelativistic potential models exploiting lattice QCD singlet free energies as potentials in the Schrödinger equation for heavy quarkonia. This agreement with previous results (see, e.g., Ref. [7]) includes also higher quarkonia resonances for which the stabilization effects is more pronounced.

5 Acknowledgments

JJ received financial support from the Bogoliubov-Infeld program for his participation in the Helmholtz International Summer Schools in Dubna. The work of DB has been supported in part by the Polish Ministry for Science and Higher Education under grant No. N N 202 0953 33 and by RFBR grant No. 08-02-01003-a.

References

- [1] T. Matsui and H. Satz, Phys. Lett. B **178** 416 (1986).
- [2] N. F. Mott, Rev. Mod. Phys. **40** 677 (1968).
- [3] R. Rapp, D. Blaschke and P. Crochet, arXiv:0807.2470 [hep-ph] (2008).
- [4] V. V. Dixit, Mod. Phys. Lett. A **5** 227 (1990).
- [5] D. Cabrera and R. Rapp, Phys. Rev. D **76** 114506 (2007);
C. Y. Wong, Phys. Rev. C **72** 034906 (2005).
- [6] M. Asakawa, T. Hatsuda and Y. Nakahara, Prog. Part. Nucl. Phys. **46** 459 (2001);
A. Jakovac, P. Petreczky, K. Petrov and A. Velytsky, Phys. Rev. D **75** 014506 (2007).
- [7] H. Satz, J. Phys. G **32** R25 (2006).
- [8] S. P. Klevansky, Rev. Mod. Phys. **64** (1992) 649.
- [9] C. Ratti, M. A. Thaler and W. Weise, Phys. Rev. D **73** 014019 (2006).
- [10] H. Hansen *et al.*, Phys. Rev. D **75** 065004 (2007).
- [11] J. I. Kapusta and C. Gale, *Finite-Temperature Field Theory*, Cambridge University Press (2006).
- [12] A. Beraudo, J. P. Blaizot and C. Ratti, Nucl. Phys. A **806** 312 (2008).
- [13] D. Blaschke, M. Buballa, A. E. Radzhabov and M. K. Volkov, Yad. Fiz. **71**, 2012 (2008).
- [14] F. Arleo, P. B. Gossiaux, T. Gousset and J. Aichelin, Phys. Rev. D **65** 014005 (2002).